

REVIEW

Apathy as Non-Motor Manifestation in Amyotrophic Lateral Sclerosis

Anca Moțățăianu¹, Ioana Ormenișan^{2*}, Rodica Bălașa¹

1. Neurology Department, George Emil Palade University of Medicine, Pharmacy, Science, and Technology of Targu Mures, Romania

2. Neurology I Clinic, County Emergency Clinical Hospital of Targu Mures, Romania

Although the term of amyotrophic lateral sclerosis (ALS) is often used interchangeably with motor neuron disease, ALS is universally accepted as a multisystem disorder. Cognitive impairment is an acknowledged feature of ALS, affecting language, memory and behaviour, and apathy is considered to be the prevalent behavioural alteration in ALS. It can be divided in three subtypes: executive, emotional and initiation apathy. Out of the three subtypes, initiation apathy is common among patients with ALS. Even in patients that do not meet the criteria for ALS with frontotemporal dementia, low-key neuropsychiatric and cognitive changes can be observed. Apathy has also been found to be systematically associated with disruptions in medial frontal cortex and subcortical structures in several neuroimaging studies that confirm the pre-existing brain lesions in the early stages of this disease. Thus, there is a growing body of evidence that motor signs and symptoms are accompanied or even preceded by cognitive and behavioural alterations, and screening for non-motor signs and symptoms can be clinically relevant.

Keywords: amyotrophic lateral sclerosis, non-motor manifestation, apathetic syndrome, depression

Received 26 October 2021 / Accepted 18 November 2021

Introduction

Amyotrophic Lateral Sclerosis (ALS) is an incurable relentlessly progressive neurodegenerative disorder causing muscle weakness and culminating in fatal dysphagia and respiratory failure [1]. It is no longer believed to be a pure motor system degeneration. Cognitive impairment is an acknowledged feature of ALS and it affects emotion processing, social cognition, executive functions and behaviour. They are all identified and described as extra-motor signs and symptoms of the disease [2,3].

Screening for Non-Motor Manifestations

According to several recent studies, more than 50% of patients with ALS experience cognitive dysfunctions over the course of the disease (i.e., changes in language, behaviour, memory, social cognition) [4]. About 30%-40% of patients with cognitive dysfunctions present with cognitive impairment alone, 10% have only behavioural impairment, and 10-15% reach criteria for frontotemporal dementia (FTD) [5].

In order to quantify these non-motor manifestations a new assessment instrument was developed by Abrahams et al. (2014) - the Edinburgh Cognitive and Behavioural ALS Screen (ECAS), a screening test developed specifically for ALS patients. Its design allows assessing cognitive performance irrespective of physical disability, which may be a confounder in ALS [6,7]. ECAS is considered the gold standard for the assessment of cognition and behaviour in ALS and it is widely used as a screening tool for ALS-specific (executive functions, verbal fluency, and language tests) and ALS non-specific subtests (memory and visuo-

spatial tests) [8], and a carer behaviour screen of five domains pertaining to FTD [7].

The Apathetic Syndrome

Apathy is acknowledged as the prevalent behavioural alteration in ALS and other neurodegenerative diseases[9]. Studies show that it occurs in 30–60% of ALS patients[10]. Marin (1996) defines apathy as “lack of motivation or decreased motivation towards goal-directed behaviours” [11]. He also refers to the symptom of apathy as loss of motivation as a result of disruption in intellect, emotion, or level of consciousness [12], suggesting thus its multidomain structure.

Strong et al. (2017) revised the criteria for ALS with behavioural impairment and included apathy as a main symptom [13]. Apathy was conventionally measured as a unidimensional symptom [10]. Nevertheless, recent findings suggest that apathy in ALS can be divided in three subtypes that can be measured using a specifically designed instrument, the Dimensional Apathy Scale (DAS): (1) executive apathy, referring to planning, attention or organization impairments, (2) emotional apathy, concerning emotion integration deficits, and (3) initiation apathy, pertaining to impaired self-generation of behaviours or cognition [10,14].

Initiation apathy, which is associated with impaired origination of thoughts or actions, has been found to be the most common subtype in patients with ALS. It is also associated with poor results on verbal fluency tests [15], which have been employed as highly sensitive instruments to measure executive dysfunction in clinical settings [3]. Thus test results show patients are lacking intrinsic response generation, planning and goal management strategies [16].

* Correspondence to: Ioana Ormenișan
E-mail: ioana.ormenisan@gmail.com

Apathy has been found to be systematically associated with disruptions in medial frontal cortex and subcortical structures in several neuroimaging studies [17]. Femiano et al. (2018) aimed to prove a potential association between brain microstructural damage and apathy in the early stages of ALS. They used diffusion tensor imaging (DTI), which is able to investigate white matter integrity, and King's clinical staging system, a validated staging system for ALS. Their findings revealed that, in early stages of disease, ALS patients without cognitive or behaviour dysfunctions may present with microstructural changes in motor and respectively, extra-motor white matter tracts, namely corticospinal tracts, midbody of corpus callosum and bilateral anterior cingulate bundles [9].

Woolley et al. (2011) aimed to determine whether apathy in ALS relates to structural changes associated with the degenerative process, also using DTI with fractional anisotropy. Their results are consistent with the previous study, confirming that there are neuroanatomical grounds for apathy in non-demented ALS patients [18]. The anterior cingulum is known to be involved in both the regulation of goal-directed and socioemotional behaviours and inhibitory control, explaining thus the strong association between the altered anterior cingulum bundle and apathy in ALS [9,19].

The inferior parietal lobule is another region believed to be involved in movement intention and movement perception, and thus disruptions in this area may be related to the occurrence of apathy [20]. Tumati et al. (2018) concluded that altered neural input from this region may affect intentional goal-directed actions. Patients find it difficult to initiate behaviour and they need instructions to perform regular or habitual actions [21].

Apathy and Depression

Depression is considered to be a marker of global neurodegeneration, albeit a nonspecific one [22]. Nevertheless, depression in patients with ALS may be an underdiagnosed comorbidity, mainly because clinical signs of cognitive impairment may confound depressive symptoms [23,24].

Caga et al. (2016) examined the relationship between apathy levels, mortality and survival time. They also assessed the link between specific symptom clusters of depression and apathy. Depression symptoms were categorised as follows: depression with demoralization and depression with anhedonia. Their results showed there was no significant association between the level of apathy and depression with demoralization (feelings of worthlessness and hopelessness). In contrast, depression with anhedonia (loss of interest or pleasure) was more frequent in patients with mild and moderate to severe apathy [25].

Differentiating between apathy and anhedonia is challenging. Abnormalities in the anterior cingulate circuits are thought to be implicated in the pathophysiology of both apathy and depression [25]. Another study identifies the ventromedial sectors of the prefrontal cortex as the neural

pathways underlying depression [26], nevertheless lesions in the dorsolateral sector of the prefrontal cortex remain a common structural marker for both apathy and depression [25].

Apathy as a Premorbid Condition

As mentioned before, recent findings suggest that patients with ALS, who do not make up for the 10 to 15% diagnosed with FTD, also present with low-key neuropsychiatric and cognitive changes [27], apathy being most often reported. Mioshi et al. (2014) study results confirm the hypothesis that neuropsychiatric symptoms precede classic motor features of ALS [27]. In another study, they also prove that significant cortical atrophy is present in patients with ALS and neuropsychiatric symptoms. However, this cortical alteration lacks in those who only exhibit motor symptoms [28]. Their results support the conclusions of the abovementioned studies of Femiano and Woolley that confirm the pre-existing brain lesions correlated with apathy in the early stages of the disease.

Interestingly enough, in the ALS context, apathy used to be consistently misread as patients being unusually compliant [6]. Mehl et al. (2017) surveyed the physicians' assessment of the personality structure of their ALS patients. The authors' endeavour was motivated by several studies and reviews on ALS where physicians described patients as "genuinely pleasant", "pleasant and warm", or "unusually stoic". They used a multidimensional personality questionnaire (the shortened version of the "NEO Five-Factor Inventory"), and they concluded that helpfulness and compliance seem to be prevalent in ALS patients' perceived behaviour [29].

Grossman et al. (2006) conducted a caregiver-based study where ALS patients were rated a lower score on "openness" as a premorbid personality trait compared to patients with other chronic medical conditions (lung cancer, multiple sclerosis or brain glioma). Openness describes individuals with conventional values, who would follow the advice of their physicians and comply with medical treatment. Their results suggest that there may be premorbid personality features that are common among patients with ALS [30].

Conclusions

Although the term of amyotrophic lateral sclerosis is often used interchangeably with motor neuron disease, ALS is universally accepted as a multisystem disorder. There is a growing body of evidence that motor signs and symptoms are accompanied or even preceded by cognitive and behavioural changes. Notwithstanding the inexorable progress and fatal outcome of ALS, screening for non-motor signs and symptoms can be clinically relevant, allowing optimal clinical approach and treatment tailored to the needs of each patient.

Given the impact of being diagnosed with an incurable disease, is there a chance that behavioural alterations be

erroneously considered a reasonable response to the diagnosis? Do lack of motivation and loss of interest manifest as a consequence of such diagnosis or are they actually a tell-tale sign of early behavioural impairment? Is apathy a prodromal phase of ALS? To what extent are the non-motor signs and symptoms concealed by physical deficits? Can we buy ALS patients more time? These are questions that remain to be addressed.

Acknowledgment

This review was supported by research grant 293/5/2020 UMFST "George Emil Palade" Targu Mures.

Authors' Contribution:

AM (Conceptualization; Supervision; Writing – review & editing)

IO (Conceptualization; Investigation; Writing – original draft)

RB (Supervision; Validation; Visualization)

References

- Cui F, Zhu W, Zhou Z, et al. Frequency and risk factor analysis of cognitive and anxiety-depressive disorders in patients with amyotrophic lateral sclerosis/motor neuron disease. *Neuropsychiatr Dis Treat*. 2015 Nov 3;11:2847-54
- Benbrika S, Desgranges B, Eustache F, Viader F. Cognitive, Emotional and Psychological Manifestations in Amyotrophic Lateral Sclerosis at Baseline and Overtime: A Review. *Front Neurosci*. 2019 Sep 10;13:951
- Abrahams S, Leigh PN, Harvey A, Vythelingum GN, Gris e D, Goldstein LH. Verbal fluency and executive dysfunction in amyotrophic lateral sclerosis (ALS). *Neuropsychologia*. 2000;38(6):734-47
- Goldstein LH, Abrahams S. Changes in cognition and behaviour in amyotrophic lateral sclerosis: nature of impairment and implications for assessment. *Lancet Neurol*. 2013 Apr;12(4):368-80
- De Marchi F, Sarnelli MF, Solara V, Bersano E, Cantello R, Mazzini L. Depression and risk of cognitive dysfunctions in amyotrophic lateral sclerosis. *Acta Neurol Scand*. 2019 May;139(5):438-445
- Pender N, Pinto-Grau M, Hardiman O. Cognitive and behavioural impairment in amyotrophic lateral sclerosis. *Curr Opin Neurol*. 2020 Oct;33(5):649-654
- Abrahams S, Newton J, Niven E, Foley J, Bak TH. Screening for cognition and behaviour changes in ALS. *Amyotroph Lateral Scler Frontotemporal Degener*. 2014 Mar;15(1-2):9-14
- Carelli L, Solca F, Faini A, et al. The Complex Interplay Between Depression/Anxiety and Executive Functioning: Insights From the ECAS in a Large ALS Population. *Front Psychol*. 2018 Apr 5;9:450
- Femiano C, Trojsi F, Caiazzo G, et al. Apathy Is Correlated with Widespread Diffusion Tensor Imaging (DTI) Impairment in Amyotrophic Lateral Sclerosis. *Behav Neurol*. 2018 Oct 22;2018:2635202
- Radakovic R, Stephenson L, Colville S, Swingler R, Chandran S, Abrahams S. Multidimensional apathy in ALS: validation of the Dimensional Apathy Scale. *J Neurol Neurosurg Psychiatry*. 2016 Jun;87(6):663-9
- Marin RS. Apathy: Concept, Syndrome, Neural Mechanisms, and Treatment. *Semin Clin Neuropsychiatry*. 1996 Oct;1(4):304-314 - abstract
- Marin RS. Apathy: a neuropsychiatric syndrome. *J Neuropsychiatry Clin Neurosci*. 1991 Summer;3(3):243-54 - abstract
- Strong MJ, Abrahams S, Goldstein LH, et al. Amyotrophic lateral sclerosis - frontotemporal spectrum disorder (ALS-FTSD): Revised diagnostic criteria. *Amyotroph Lateral Scler Frontotemporal Degener*. 2017 May;18(3-4):153-174
- Levy R. Apathy: a pathology of goal-directed behaviour: a new concept of the clinic and pathophysiology of apathy. *Rev Neurol (Paris)*. 2012 Aug-Sep;168(8-9):585-97
- Caga J, Hsieh S, Highton-Williamson E, et al. Apathy and its impact on patient outcome in amyotrophic lateral sclerosis. *J Neurol*. 2018 Jan;265(1):187-193
- Radakovic R, Stephenson L, Newton J, et al. Multidimensional apathy and executive dysfunction in amyotrophic lateral sclerosis. *Cortex*. 2017 Sep;94:142-151
- Le Heron C, Holroyd CB, Salamone J, Husain M. Brain mechanisms underlying apathy. *J Neurol Neurosurg Psychiatry*. 2019 Mar;90(3):302-312
- Woolley SC, Zhang Y, Schuff N, Weiner MW, Katz JS. Neuroanatomical correlates of apathy in ALS using 4 Tesla diffusion tensor MRI. *Amyotroph Lateral Scler*. 2011 Jan;12(1):52-8
- Massimo L, Powers JP, Evans LK, et al. Apathy in Frontotemporal Degeneration: Neuroanatomical Evidence of Impaired Goal-directed Behavior. *Front Hum Neurosci*. 2015 Nov 10;9:611
- Caga J, Tu S, Dharmadasa T, et al. Apathy is associated with parietal cortical-subcortical dysfunction in ALS. *Cortex*. 2021 Mar 23;S0010-9452(21)00103-
- Tumati S, Martens S, de Jong BM, Aleman A. Lateral parietal cortex in the generation of behavior: Implications for apathy. *Prog Neurobiol*. 2019 Apr;175:20-34
- De Marchi F, Sarnelli MF, Solara V, Bersano E, Cantello R, Mazzini L. Depression and risk of cognitive dysfunctions in amyotrophic lateral sclerosis. *Acta Neurol Scand*. 2019 May;139(5):438-445
- Motataianu A, Andone S, Radu C, et al. Predictors of Depression in Caucasian Patients with Amyotrophic Lateral Sclerosis in Romania. *Brain Sci*. 2020 Jul 22;10(8):470
- K rner S, Kollwe K, Ilsemann J, et al. Prevalence and prognostic impact of comorbidities in amyotrophic lateral sclerosis. *Eur J Neurol*. 2013 Apr;20(4):647-54
- Caga J, Turner MR, Hsieh S, et al. Apathy is associated with poor prognosis in amyotrophic lateral sclerosis. *Eur J Neurol*. 2016 May;23(5):891-7
- Koenigs M, Grafman J. The functional neuroanatomy of depression: distinct roles for ventromedial and dorsolateral prefrontal cortex. *Behav Brain Res*. 2009 Aug 12;201(2):239-43
- Mioshi E, Caga J, Lillo P, et al. Neuropsychiatric changes precede classic motor symptoms in ALS and do not affect survival. *Neurology*. 2014 Jan 14;82(2):149-55
- Mioshi E, Lillo P, Yew B, et al. Cortical atrophy in ALS is critically associated with neuropsychiatric and cognitive changes. *Neurology*. 2013 Mar 19;80(12):1117-23
- Mehl T, Jordan B, Zierz S. "Patients with amyotrophic lateral sclerosis (ALS) are usually nice persons"-How physicians experienced in ALS see the personality characteristics of their patients. *Brain Behav*. 2016 Nov 10;7(1):e00599
- Grossman AB, Levin BE, Bradley WG. Premorbid personality characteristics of patients with ALS. *Amyotroph Lateral Scler*. 2006 Mar;7(1):27-31